



**SINA**

Conference 2015 Tulln/Donau

Stable Isotope Network Austria

**14<sup>th</sup> Austrian**

**Stable Isotope**

**Network Meeting**

**Tulln/Donau, November 26<sup>th</sup>-27<sup>th</sup> 2015**

**Austrian Institute of Technology GmbH  
Health and Environment Department**

**University of Natural Resources and Life Sciences  
Department of Chemistry (VIRIS Lab)**

Austrian Institute of Technology and University of  
Natural Resources and Life Sciences

## **14<sup>th</sup> Stable Isotope Network Austria Meeting**

Tulln/Donau 26<sup>th</sup>-27<sup>th</sup> November 2015

### **PROGRAM AND ABSTRACT VOLUME**

Editorial: STEFAN WYHLIDAL, ANDREA WATZINGER, REBECCA HOOD-NOWOTNY,  
THOMAS PROHASKA

To be cited as Wyhlidal, S., Watzinger, A., Hood-Nowotny, R., Prohaska, T., (eds): 14<sup>th</sup> Austrian Stable  
isotope user group meeting. Austrian Institute of Technology, 39p., Tulln/Donau, 2015

Impressum:

Alle Rechte für das In- und Ausland vorbehalten.

Copyright: Austrian Institute of Technology, Konrad-Lorenz-Str. 24, 3430 Tulln, Austria  
Medieninhaber, Herausgeber und Verleger: Austrian Institute of Technology, Tulln/Donau  
homepage: [www.ait.ac.at](http://www.ait.ac.at)

## 21 Poster Presentation

### TOWARDS A SR ISOSCAPE OF AUSTRIA FOR THE DETERMINATION OF GROWTH REGIONS OF PREHISTORIC WOOD

Horsky, M.<sup>1</sup>, Tintner, J.<sup>2</sup>, Grabner, M.<sup>2</sup>, Kowarik, K.<sup>3</sup>, Reschreiter, H.<sup>3</sup>,  
Teufl, K.<sup>4</sup>, Horacek, M.<sup>5</sup> & Prohaska, T.<sup>1</sup>

<sup>1</sup>University of Natural Resources and Life Sciences, Vienna, Department of Chemistry, Division of Analytical Chemistry, VIRIS Laboratory, Konrad-Lorenz-Str. 24, 3430 Tulln, Austria

<sup>2</sup>University of Natural Resources and Life Sciences, Vienna, Department of Material Sciences and Process Engineering, Institute of Wood Science and Technology, Konrad-Lorenz-Str. 24, 3430 Tulln, Austria

<sup>3</sup>Natural History Museum Vienna, Department of Prehistory, Burgring 7, 1010 Vienna, Austria

<sup>4</sup>Josephinum Research, Rottenhauser Straße 1, 3250 Wieselburg, Austria

<sup>5</sup>BLT Wieselburg, Lehr- und Forschungszentrum Francisco-Josephinum, Rottenhauser Straße 1, 3250 Wieselburg, Austria

e-mail:monika.horsky@boku.ac.at

Wood artefacts from a prehistoric salt mine in Hallstatt, Austria, present a unique archive of information on Bronze and Iron Age mining. For instance, the geochemical signatures of growth locations are stored in the finds, though masked by contaminating minerals attributable to the storage conditions. Trade is assumed for certain archaeological finds. Consequently, analysis of the radiogenic strontium isotope amount ratio  $n(^{87}\text{Sr})/n(^{86}\text{Sr})$  has been applied to investigate the geographic origin of these artefacts, in order to allow conclusions on trade routes.

In order to reveal the biogenic signatures of the prehistoric finds, a decontamination method based on acid leaching was developed. It enabled the separation of biogenic from diagenetic Sr. A mixing model was adopted to account for possibly incomplete removal of the latter. In addition to Hallstatt, seven regions in Austria were selected for sampling of modern trees based on known settlements in the time period of interest. The geological bedrock variability was considered within all regions for the definition of sampling spots, which resulted in a total of 26 locations. Drill cores from four tree species represented in the archaeological finds (i.e. *Picea abies*, *Abies alba*, *Fagus sylvatica* and *Quercus* sp.) were sampled upon availability. Sr isotope ratios were measured in wood digests after Sr/matrix separation using multicollector inductively coupled plasma-mass spectrometry (MC ICP-MS).

The isotopic signature of bioavailable Sr that was obtained from modern trees reflects the geological heterogeneity in Austria, which challenges the creation of an isoscape and its applicability to distinct provenance determination. Different bedrock types can be distinguished by their  $n(^{87}\text{Sr})/n(^{86}\text{Sr})$ . Furthermore, the data indicate that the isotope ratios of bio-available Sr within one geological substrate also vary strongly. The results highlight the importance to consider even small scale geological and environmental variability in a comprehensive sampling strategy for a reliable application of Sr isotope ratio analysis to the determination of origin of biogenic material.

This contribution was also presented at the DocDay 2015, Tulln, Austria.